arXiv:1203.5487v1 [math.GT] 25 Mar 2012

ON UNSTABILZED GENUS THREE CRITICAL HEEGAARD SURFACES

JUNGSOO KIM

Abstract. Let M be a compact orientable irreducible 3-manifold and H be an unstabilized genus three Heegaard splitting of M. In this article, we will define a simplicial complex of weak reducing pairs for H and find several properties of this complex. Using this method, we will prove that an unstabilized Heegaard splitting of genus three is critical in a certain condition. In addition, we will show that the standard genus three Heegaard splitting for T^3 and the induced Heegaard splitting of the three component chain exterior by a certain tunnel system are critical as examples of the main theorem.

Contents

1. Introduction and result 1
2. Critical surfaces and the complex of weak reducing pairs 3
3. The proof of Theorem 1.1 7
4. The proof of Corollary 1.2 17
5. Examples of Theorem 1.1 20
5.1. The three-torus T^3 20
5.2. The three component chain exterior 21
Acknowledgments 22
References 22

1. Introduction and result

Throughout this paper, all surfaces and 3-manifolds will be taken to be compact and orientable. In [1], Bachman introduced the concept *"a critical surface"* and he proved several theorems about incompressible surfaces, the number of Heegaard splittings with respect to its genus, and the minimal genus common stabilization. Since a critical surface has disjoint compressions on it's both sides, if the surface is a Heegaard surface, then the splitting is weakly reducible, i.e. a critical Heegaard splitting is a kind of weakly reducible splitting. But in some aspects, it shares common properties with strongly irreducible splittings. For example, if the splitting is strongly irreducible or critical, then the manifold is irreducible (Lemma 3.5 of [2].) Indeed, the intersection of an incompressible surface S and a Heegaard surface F can be isotoped essential on both S and F if the splitting is critical or strongly irreducible (see Theorem 5.1 of [1] and Lemma 6 of [12].) Bachman also proved

---

Date: 25, Mar, 2012.

2000 *Mathematics Subject Classification.* 57M50.

Gordon's conjecture by using the series of generalized Heegaard splittings and critical Heegaard splittings (see [2].) In his recent work [3], he also introduced the concept *"topologically minimal surfaces"*, where a strongly irreducible surface is an index 1 topological minimal surface, and a critical surface is an index 2 topological minimal surface, this is a way to regard strongly irreducible surfaces and critical surfaces in a unified viewpoint.

Although critical Heegaard splittings have many powerful properties as proved in Bachman's recent works, it is not easy to determine whether a weakly reducible splitting is critical. For genus two 3-manifolds, a weakly reducible splitting is also a reducible splitting (see [14], the proof can be extended easily to the case with non-empty boundary.) Moreover, if a 3-manifold has a reducible splitting, then the manifold is reducible or the splitting is stabilized (see [9].) Since a reducible manifold cannot have a critical Heegaard splitting, we need to consider the manifolds of genus at least three. In this article, we will prove Theorem 1.1.

Theorem 1.1. *Let M be an orientable irreducible 3-manifold and $H = (V, W; F)$ be an unstabilized genus three Heegaard splitting. Suppose that there is no weak reducing pair such that each of both disks cuts off a solid torus in its compression body. If we can choose two weak reducing pairs (D_0, E_0) and (D_1, E_1) such that $\partial D_0 \cap \partial E_1 \neq \emptyset$ and $\partial D_1 \cap \partial E_0 \neq \emptyset$ up to isotopy, then H is critical.*

We can induce following corollary,

Corollary 1.2. *Let M be a closed orientable irreducible 3-manifold and $H = (V, W; F)$ be an unstabilized genus three Heegaard splitting. Suppose that H is not an amalgamation of two genus two splittings along a torus. If we can choose two weak reducing pairs (D_0, E_0) and (D_1, E_1) such that $\partial D_0 \cap \partial E_1 \neq \emptyset$ and $\partial D_1 \cap \partial E_0 \neq \emptyset$ up to isotopy, then H is critical.*

In 2002, Moriah proved that if an orientable 3-manifold has a weakly reducible Heegaard splitting of minimal genus, then M contains an essential surface of positive genus (see [8].) (In 1987, Casson and Gordon proved this when M is closed and the Heegaard splitting is irreducible (see [4].)) By using Bachman's result (Theorem 5.1 of [1]), we can directly get the following corollary.

Corollary 1.3. *Let M be an orientable irreducible genus three 3-manifold and $H = (V, W; F)$ be a minimal genus Heegaard splitting. Suppose that there is no weak reducing pair such that each of both disks cuts off a solid torus in its compression body. If we can choose two weak reducing pairs (D_0, E_0) and (D_1, E_1) such that $\partial D_0 \cap \partial E_1 \neq \emptyset$ and $\partial D_1 \cap \partial E_0 \neq \emptyset$ up to isotopy, then there is an essential surface S of positive genus in M such that $F \cap S$ is essential on both F and S.*

In a genus three Heegaard splitting, if each of both disks for a weak reducing pair cuts off a solid torus in its compression body, then both disks are separating in their compression bodies. Therefore, we can use Theorem 1.1 and its corollaries when if there is no weak reducing pair whose two disks are separating in their compression bodies.

This article is organized as follows. In section 2, we introduce some basic notions, define the complex of weak reducing pairs $\mathcal{WR}$, and find basic properties of $\mathcal{WR}$, especially about the criticality of a Heegaard surface, the shapes of simplices in $\mathcal{WR}$, and stabilizations of a Heegaard surface. In section 3, we will induce special properties of $\mathcal{WR}$ when the manifold is irreducible, the splitting is unstabilzed, and the genus is three. By using these properties, we will prove Theorem 1.1. In section 4, we will find additional properties of $\mathcal{WR}$ when the manifold is closed and prove Corollary 1.2. In section 5, we will show that the standard splitting of genus three for the three-torus T^3 is critical and find an incompressible surface S' which satisfies Corollary 1.3 as an example of Theorem 1.1. In addition, we will show that the induced Heegaard splitting of the three component chain exterior by a certain tunnel system is critical.

2. Critical surfaces and the complex of weak reducing pairs

A *compression body* is a 3-manifold which can be obtained by starting with some closed, orientable, connected surface F, forming the product $F \times I$, attaching some number of 2-handles to $F \times \{1\}$ and capping off all resulting 2-sphere boundary components that are not contained in $F \times \{0\}$ with 3-balls. The boundary component $F \times \{0\}$ is referred to as ∂_+. The rest of the boundary is referred to as ∂_-. A *Heegaard splitting* of a 3-manifold M is an expression of M as a union $V \cup_F W$, where V and W are compression bodies that intersect in a transversally oriented surface $F = \partial_+ V = \partial_+ W$. We will use the expression $(V, W; F)$ for a Heegaard splitting. If $(V, W; F)$ is a Heegaard splitting of M then we say that F is a *Heegaard surface*. We say that the pair (D, E) is a *weak reducing pair* for F if $D \subset V$ and $E \subset W$ are disjoint compressing disks. A Heegaard surface is *strongly irreducible* if it is compressible to both sides but has no weak reducing pairs. From now, we will use the letter "D" for compressing disks in V, "E" for compressing disks in W, "F" for the Heegaard surface in the given Heegaard splitting, and "H" for the name of a Heegaard splitting.

Definition 2.1 (D. Bachman, Definition 3.3 of [2])**.** Let F be a Heegaard surface in some 3-manifold which is compressible to both sides. The surface F is *critical* if the set of all compressing disks for F can be partitioned into subsets C_0 and C_1 such that the follows hold.

(1) For each $i = 0, 1$ there is at least one weak reducing pair (D_i, E_i), where D_i, $E_i \in C_i$.
(2) If $D \in C_i$ and $E \in C_j$, then (D, E) is not a weak reducing pair for $i \neq j$.

Note that the definition of *"critical surface"* of [2] is significantly simpler and slightly weaker, than the one given in [1]. In other words, anything that was considered critical in [1] is considered critical here as well.

Let M be a compact orientable 3-manifold with possibly non-empty boundary, and suppose that M has a weakly reducible Heegaard splitting $H = (V, W; F)$.

Let D and D' be compressing disks in V. By abuse of terminology, we will just say that $D \cap D' = \emptyset$ ($E \cap E' = \emptyset$ resp.) if they are not isotopic in V (W resp.) and D misses D' (E misses E' resp.). In the case D and D' (E and E' resp.) are isotopic in V (W resp.), we will denote it as $D = D'$ ($E = E'$ resp.) even if D misses D' (E misses E' resp.) Similarly, we will just say that $\partial D \cap \partial E = \emptyset$ if ∂D and ∂E are not isotopic in F and ∂D misses ∂E. In the case ∂D and ∂E are isotopic in F, we will denote it as $\partial D = \partial E$ even if ∂D misses ∂E. Therefore, for a weak reducing pair (D, E), we get either $\partial D \cap \partial E = \emptyset$ or $\partial D = \partial E$. If there is a weak reducing pair which holds the latter case, the splitting is reducible.

Definition 2.2. Define *the complex of weak reducing pairs* $\mathcal{WR}$ as follows.

(1) Each vertex of $\mathcal{WR}$ is a weak reducing pair for F.

(2) Two vertices $v = (D, E)$ and $w = (D', E')$ in $\mathcal{WR}$ are the same if and only if $D = D'$ and $E = E'$.
(3) Assign an edge (or a 1-simplex) e between two different vertices $v = (D, E)$ and $w = (D', E') \in \mathcal{WR}$ if (a) $D = D'$ and $E \cap E' = \emptyset$, or (b) $D \cap D' = \emptyset$, and $E = E'$. The edge e between v and w is determined uniquely since it is impossible that (a) and (b) occur simultaneously.

We define an *n-simplex* in $\mathcal{WR}$ by $(n+1)$-vertices $v_0, \cdots, v_n$ if there is an edge between v_i and v_j for every choice of i and j, where $0 \leq i \neq j \leq n$. $\mathcal{WR}$ depends on the manifold and the Heegaard splitting. If we need to make sure of the Heegaard surface or the manifold, denote the complex as $\mathcal{WR}(F)$ or $\mathcal{WR}(M; F)$.

It may be possible that an edge $(D_1, E) - (D_2, E)$ exists but $\partial D_1 = \partial D_2$ if the manifold has a sphere boundary. But if we consider irreducible manifolds other than B^3, then a Heegaard splitting consists of non-punctured compression bodies. Therefore, if M is an irreducible manifold other than B^3 and $\partial D_1 = \partial D_2$, then we can isotope D_1 and D_2 in V so that $D_1 = D_2$ since a compression body is irreducible. Similary, if M is an irreducible manifold other than B^3, $\partial D_1 \cap \partial D_2 = \emptyset$, and ∂D_1 is not isotopic to ∂D_2 in F, then we can isotope D_1 and D_2 in V so that $D_1 \cap D_2 = \emptyset$.

In section 8 of [2], D. Bachman considered a sequence of compressing disks

$$D = D_0 - E = E_0 - D_1 - E_1 - \cdots - D' = D_m - E' = E_m,$$

where (a) $D_i = D_{i+1}$ or $D_i \cap D_{i+1} = \emptyset$ and (b) $E_i = E_{i+1}$ or $E_i \cap E_{i+1} = \emptyset$ for each $0 \leq i \leq m-1$ and both (D_i, E_i) for $0 \leq i \leq m$ and (D_{i+1}, E_i) for $0 \leq i \leq m-1$ are weak reducing pairs. He defined the distance between two weak reducing pairs (D, E) and (D', E') using the minimal length of this sequence. In particular, if there is no such a sequence between them, the distance is defined as ∞. He also proved that if the distance between two weak reducing pairs is ∞, then the Heegaard surface is critical. We can rewrite this result in terms of $\mathcal{WR}$ as follows.

Proposition 2.3 (D. Bachman, Lemma 8.5 of [2])**.** *If $\mathcal{WR}$ is disconnected, then the Heegaard surface is critical.*

Proof. We will prove that two vertices (D, E) and (D', E') in $\mathcal{WR}$ are connected by a union of 1-simplices if and only if the distance between (D, E) and (D', E') is finite. Therefore, this is equivalent to Lemma 8.5 of [2].

Suppose that (D, E) and (D', E') in $\mathcal{WR}$ are connected by a union of m 1-simplices,

$$(D = D_0, E = E_0) -_{e_1} (D_1, E_1) -_{e_2} \cdots -_{e_m} (D' = D_m, E' = E_m).$$

Each 1-simplex $(D_{i-1}, E_{i-1}) -_{e_i} (D_i, E_i)$ denotes that (a) $D_{i-1} = D_i$ and $E_{i-1} \cap E_i = \emptyset$ or (b) $D_{i-1} \cap D_i = \emptyset$ and $E_{i-1} = E_i$. In both cases, we can assign a sequence like following to e_i,

$$D_{i-1} - E_{i-1} - D_{i-1} - E_i - D_i - E_i.$$

Similarly, for the 1-simplex $(D_i, E_i) -_{e_{i+1}} (D_{i+1}, E_{i+1})$, we can assign a sequence like following to e_{i+1},

$$D_i - E_i - D_i - E_{i+1} - D_{i+1} - E_{i+1}.$$

The sequence corresponding to e_i ends with $D_i - E_i$ and that corresponding to e_{i+1} starts with $D_i - E_i$. Therefore, if we connect these two sequences identifying the common segment $D_i - E_i$ for all $1 \leq i \leq m-1$, then we get the wanted sequence, i.e. the distance between (D, E) and (D', E') is finite.

Conversely, suppose that the distance between (D, E) and (D', E') is finite, i.e. there is a sequence like following,

$$D = D_0 - E = E_0 - D_1 - E_1 - \cdots - D' = D_m - E' = E_m.$$

Now we will follow the procedure described below.

1. Read the sequence from $D_0 - E_0$. Initially, we assign a vertex (D_0, E_0).
2. For $D_0 - E_0 - D_1$, if $D_0 \cap D_1 = \emptyset$, then we assign a 1-simplex e in $\mathcal{WR}$, $(D_0, E_0) -_e (D_1, E_0)$. If $D_0 = D_1$, then we define (D_1, E_0) as (D_0, E_0).
3. For $E_0 - D_1 - E_1$, if $E_0 \cap E_1 = \emptyset$, then we assign a 1-simplex f in $\mathcal{WR}$, $(D_1, E_0) -_f (D_1, E_1)$. If $E_0 = E_1$, then we define (D_1, E_1) as (D_1, E_0).
4. We repeat the steps (2) and (3), i.e. consider $D_i - E_i - D_{i+1}$ and $E_i - D_{i+1} - E_{i+1}$ for all $1 \leq i \leq m-1$.
5. Finish after the step $E_{m-1} - D_m - E_m$.

Here, we get a union of 1-simplices which connects (D, E) and (D', E') in $\mathcal{WR}$. □

In terms of $\mathcal{WR}$, we can also get the following lemma.

Lemma 2.4 (D. Bachman, Lemma 8.4 of [2])**.** *If there are two different vertices $v = (D, E)$ and $w = (D, E')$, or $v = (D, E)$ and $w = (D', E)$ in $\mathcal{WR}$, then there is a path from v to w in $\mathcal{WR}$.*

We denote a 2-simplex Δ determined by the three vertices u, v, w in $\mathcal{WR}$ as $u -_e v -_f w -_g u$, where e is the 1-simplex in Δ between u and v, f is that between v and w, and g is that between w and u.

Lemma 2.5. *Suppose that Δ is a 2-simplex in $\mathcal{WR}$ determined by three vertices u, v, w. Then Δ has the form $(D, E) -_e (D, E') -_f (D, E'') -_g (D, E)$ or $(D, E) -_e (D', E) -_f (D'', E) -_g (D, E)$.*

Proof. If Δ does not have the form $(D, E) -_e (D, E') -_f (D, E'') -_g (D, E)$ or $(D, E) -_e (D', E) -_f (D'', E) -_g (D, E)$, then we get two cases by fixing the initial and terminal vertices as (D, E) and reading $u -_e v -_f w -_g u$ from the left.

Case 1. $\Delta = (D, E) -_e (D, E') -_f (D', E') -_g (D, E)$.
From the edge e, $E \cap E' = \emptyset$. Therefore, we get $D = D'$ from the edge g, this contradicts the existence of the edge f.

Case 2. $\Delta = (D, E) -_e (D', E) -_f (D', E') -_g (D, E)$.
From the edge e, $D \cap D' = \emptyset$. Therefore, we get $E = E'$ from the edge g, this contradicts the existence of the edge f. □

Lemma 2.6. *Suppose that Σ is an n-simplex in $\mathcal{WR}$ determined by $v_0, \cdots, v_n$. Then the vertices of Σ have the form (a) $v_0 = (D_0, E)$, $\cdots$ $v_n = (D_n, E)$ or (b) $v_0 = (D, E_0), \cdots, v_n = (D, E_n)$.*

Proof. We will use an induction argument for n. For $n = 2$, we already proved in Lemma 2.5. Suppose that the statement of Lemma 2.6 holds for $n = k \geq 2$, and consider a $(k+1)$-simplex Σ determined by $v_0, \cdots, v_{k+1}$. Let Δ be the k-subsimplex of Σ determined by $v_0, \cdots, v_k$, and Δ' be the k-subsimplex of Σ determined by $v_0, \cdots, v_{k-1}, v_{k+1}$. If we consider Δ, then we get the following two cases.



FIGURE 1. We can take a weak reducing pair (D, E) such that $D = \bar{D}$.

(1) the vertices of Δ have the form $v_0 = (D_0, E)$, $\cdots$, $v_k = (D_k, E)$, or
(2) the vertices of Δ have the form $v_0 = (D, E_0)$, $\cdots$, $v_k = (D, E_k)$.

In the case (1), the vertices of Δ' have the form

$$v_0 = (D_0, E), \cdots, v_{k-1} = (D_{k-1}, E), v_{k+1}.$$

Since $k \geq 2$, we can assume the existence of $v_1 = (D_1, E)$. By the induction hypothesis, v_{k+1} must have the form (D_{k+1}, E). Similarly, in the case (2), the vertices of Δ' have the form

$$v_0 = (D, E_0), \cdots, v_{k-1} = (D, E_{k-1}), v_{k+1} = (D, E_{k+1}).$$

Therefore, the statement holds when $n = k + 1$. □

Let $\bar{D} \subset V$, $\bar{E} \subset W$ be compressing disks, where $\partial\bar{D}$ and $\partial\bar{E}$ intersect transversely in a single point. Such a pair of disks is called a *canceling pair* of disks for the splitting. If there is a canceling pair, then we call the splitting *stabilized*.

Let F be a surface and D be a compressing disk of F. If we compress F along D, we call the compressed surface *"F_D"*. Lemma 2.7 gives a connection of a vertex in $\mathcal{WR}$ with a stabilized Heegaard splitting.

Lemma 2.7. *If $\{\bar{D}, \bar{E}\}$ is a canceling pair of a Heegaard splitting $H = (V, W; F)$ and the genus of F is at least two. Then, there is a weak reducing pair (D, E) such that $D = \bar{D}$ or $E = \bar{E}$.*

Proof. Let C the boundary of a regular neighborhood of $\partial\bar{D} \cup \partial\bar{E}$ in F. Then there is a simple closed curve C' in F in the same isotopy class of C in F, such that C' bounds a disk in $F_{\bar{E}}$ ($F_{\bar{D}}$ resp.). If we push the interior of this disk slightly into W (V resp), we get a disk E (D resp), such that $\bar{D} \cap E = \emptyset$ ($\bar{E} \cap D = \emptyset$ resp). If we choose $D = \bar{D}$ ($E = \bar{E}$ resp), then we get the wanted weak reducing pair (D, E) (see Figure 1.) □

Definition 2.8. Let $v = (D, E)$ be a vertex in $\mathcal{WR}$.

(1) If there is a canceling pair $\{\bar{D}, \bar{E}\}$ such that $D = \bar{D}$, then we replace F by F_D, and redefine $\mathcal{WR}(F)$ as $\mathcal{WR}(F_D)$.
(2) If there is a canceling pair $\{\bar{D}, \bar{E}\}$ such that $E = \bar{E}$, then we replace F by F_E, and redefine $\mathcal{WR}(F)$ as $\mathcal{WR}(F_E)$.

(Of course, the Heegaard splitting is also changed if one of (1) and (2) holds since the Heegaard surface is changed.) If one of (1) and (2) holds for a vertex v in $\mathcal{WR}$, we will call this procedure a *destabilization of $\mathcal{WR}$ for the vertex v*. We can repeat



FIGURE 2. If $\mathcal{WR}$ is n-stabilized, then there is an $(n-1)$-simplex determined by (D_1, E), $\cdots$, (D_n, E).

destabilizations of $\mathcal{WR}(F)$ until we cannot destabilize $\mathcal{WR}(F)$ for any vertex in $\mathcal{WR}(F)$. We call the resulting Heegaard splitting and $\mathcal{WR}(F)$ *unstabilized.* If we need n-destabilization steps to make $\mathcal{WR}$ unstabilized, we call the initial Heegaard splitting *n-stabilized* and this procedure an *n-destabilization.*

Definition 2.9. If there exists $n \in \mathbb{Z}$ such that $\dim(\Delta) \le n$ for every simplex Δ in $\mathcal{WR}$ and n is such a smallest integer, then we call *n the dimension of $\mathcal{WR}$.*

If $\mathcal{WR}$ is n-stabilized, then it is easy to see that there are n mutually disjoint non-isotopic essential disks D_1, $\cdots$, D_n in V, and an essential disk E in W s.t. $\partial E \cap \partial D_i = \emptyset$ for $1 \le i \le n$ (see Figure 2.) Therefore we get an $(n-1)$-simplex by n-vertices (D_1, E), $\cdots$, (D_n, E) in $\mathcal{WR}$, i.e. the dimension of $\mathcal{WR}$ is at least $n-1$.

3. THE PROOF OF THEOREM 1.1

In this section, we will discuss about the proof of Theorem 1.1.

Assume M, H, and F as in Theorem 1.1. Since the only irreducible 3-manifold which has a boundary component isomorphic to a 2-sphere is B^3 and a Heegaard splitting of genus three of B^3 is stabilized by using Waldhausen's Theorem (see [15]), we do not consider B^3 and assume that a boundary component of M is not homeomorphic to a 2-sphere.

Suppose that there is an m-simplex Δ in $\mathcal{WR}$. If we compress F along the disks of Δ, we get a union of surfaces. We will prove that if the splitting H is unstabilzed, then there is no 2-sphere component determined by some disks of Δ which come from different compression bodies.

Lemma 3.1. *Suppose that M is irreducible 3-manifold and $H = (V, W; F)$ is an unstabilized Heegaard splitting of M. Let Δ be an m-simplex determined by the vertices (D_0, E), $\cdots$, (D_m, E). Then, $F_{D_0\cdots D_m E}$ cannot have a S^2 component determined by E and some D_i's. Similarly, if Δ is determined by (D, E_0), $\cdots$, (D, E_m), then we get the same result for $F_{DE_0\cdots E_m}$.*

Proof. Suppose that there is a S^2 component S in $F_{D_0\cdots D_m E}$ determined by E and some D_i's. We can assume that

$$S = \bar{S} \cup (\cup \bar{D}_i) \cup (\cup \bar{E}_j),$$

where $\bar{S}$ is a planar surface, $\{\bar{D}_i\}_i$ is a set of parallel copies of the D_i's, and $\{\bar{E}_j\}_j$ is a set of parallel copies of E. Note that $\{\bar{E}_j\}_j$ consists of a disk or two disks. Let us consider a simple closed curve α in the interior of $\bar{S}$ which separates $\cup \bar{D}_i$ and

$\cup \bar{E}_j$ in S. α bounds two disks B_1 and B_2 in S, where B_1 contains $\cup \bar{D}_i$ and B_2 contains $\cup \bar{E}_j$. If we isotope the interior of B_1 and B_2 slightly, then we can assume that B_1 is properly embedded in V and B_2 is properly embedded in W. Moreover B_1 and B_2 are essential disks in their compression bodies otherwise each of the D_i's is inessential in V or E is inessential in W. Therefore, the splitting is reducible, i.e. the manifold is reducible or the splitting is stabilized (see Corollary 3.4.1 of [9].) This contradicts the hypothesis of the lemma. □

Lemma 3.2. *Suppose that M is an irreducible 3-manifold. Let $H = (V, W; F)$ be an unstabilized genus three Heegaard splitting of M. Then* $\dim(\mathcal{WR}) \leq 1$.

Proof. Let $v = (D_0, E)$ be a vertex in $\mathcal{WR}$, and without loss of generality, suppose that v is contained in an m-simplex Δ determined by $m+1$ vertices (D_0, E), $\cdots$, (D_m, E). Now ∂D_0, $\cdots$, ∂D_m, and ∂E are mutually disjoint and non-isotopic essential curves in F. (If some ∂D_i is isotopic to ∂E in F, then the splitting is reducible. But this leads that M is reducible or the splitting is stabilized.)

Case 1. ∂E is non-separating in F.

Case 1-1. $F_{D_0 \cdots D_m E}$ is connected.
This means that ∂D_0, $\cdots$, ∂D_m, and ∂E are non-separating in F. Now $F_{D_0 \cdots D_m E}$ is a surface whose genus is at most one. If $F_{D_0 \cdots D_m E}$ is a torus, then we get $m = 0$. Otherwise, $F_{D_0 \cdots D_m E}$ is a 2-sphere and $m = 1$, but this contradicts Lemma 3.1.

Case 1-2. $F_{D_0 \cdots D_m E}$ has two or more components.
In this case, there are one or two components in $F_{D_0 \cdots D_m E}$ such that each of these components is determined by E and some D_i's. Let $\bar{F}_{D_0 \cdots D_m E}$ be one of such components and suppose that E and D_0, $\cdots$, D_k, without loss of generality, determine $\bar{F}_{D_0 \cdots D_m E}$. If there are two such components, then we choose the one which makes k smallest as $\bar{F}_{D_0 \cdots D_m E}$. Assume that D_j is superfluous to determine $\bar{F}_{D_0 \cdots D_m E}$ if $j > k$. We can assume that the genus of $\bar{F}_{D_0 \cdots D_m E}$ is at least one by Lemma 3.1.

Case 1-2-1. If $k = 0$, then either $\partial E \cup \partial D_0$ cuts F into two twice-punctured tori where ∂D_0 is also non-separating in F or ∂D_0 is separating in F.

Assume that the former case. Suppose that $m \geq 1$ and let T be the twice-punctured torus component which ∂D_1 in contained in. Then, ∂D_1 is an essential simple closed curve in T which is not ∂-parallel. If ∂D_1 is separating in T, then $\partial D_0 \cup \partial D_1 \cup \partial E$ cuts off a pair of pants P from F. Hence, $F_{D_0 D_1 E}$ has a S^2 component determined by D_0, D_1, and E. If we apply Lemma 3.1 to the 1-subsimplex of Δ determined by (D_0, E) and (D_1, E), then we get a contradiction. If ∂D_1 is non-separating in T, then $F_{D_0 D_1 E}$ has a S^2 component determined by D_0, D_1, and E. Also, we get a contradiction similarly. Therefore, we get $m = 0$.

Now we assume the latter case. Let $F_{D_0} = F'_{D_0} \cup F''_{D_0}$, where $g(F'_{D_0}) = 2$ and $g(F''_{D_0}) = 1$. Suppose that $\partial E \subset F''_{D_0}$. Since ∂E is not isotopic to ∂D_0 in F, ∂E must be essential in F''_{D_0}, i.e. $F_{D_0 E}$ has a S^2-component determined by D_0 and E. If we apply Lemma 3.1 to the 0-subsimplex of Δ determined by (D_0, E), then we get a contradiction. Therefore, we get $\partial E \subset F'_{D_0}$. Moreover, $k = 0$ implies ∂D_1, $\cdots$, $\partial D_m \subset F''_{D_0}$. Since there is only one isotopy class for non-trivial disjoint simple closed curves in F''_{D_0}, we get $m \leq 1$. (We do not consider the curves isotopic to ∂D_0.) In particular, ∂D_1 must be non-separating in F if it exists.

Case 1-2-2. If $k \geq 1$ and at least two of ∂D_0, $\cdots$, ∂D_m are separating in F. Assume that ∂D_α and ∂D_β for $0 \leq \alpha \neq \beta \leq m$ are separating in F. Then,

$F_{D_\alpha D_\beta} = F'_{D_\alpha D_\beta} \cup F''_{D_\alpha D_\beta} \cup F'''_{D_\alpha D_\beta}$, where the three components of $F_{D_\alpha D_\beta}$ are genus one surfaces. Let $F'_{D_\alpha D_\beta}$ ($F'''_{D_\alpha D_\beta}$ resp.) be the one determined by only D_α (D_β resp.) and $F''_{D_\alpha D_\beta}$ be the one determined by both D_α and D_β. Suppose that $\partial E \subset F'_{D_\alpha D_\beta}$ or $F'''_{D_\alpha D_\beta}$. In this case, ∂E is essential in $F'_{D_\alpha D_\beta}$ or $F'''_{D_\alpha D_\beta}$, respectively. This means that we get a 2-sphere component in $F_{D_\alpha D_\beta E}$ determined by E and D_α, or E and D_β. Therefore, if we apply Lemma 3.1 to the 1-subsimplex of Δ determined by (D_α, E) and (D_β, E), then we get a contradiction. Now we suppose that $\partial E \subset F''_{D_\alpha D_\beta}$. We can assume that there are two disks D'_α and D'_β in $F''_{D_\alpha D_\beta}$ such that D'_α (D'_β resp.) is a parallel copy of D_α (D_β resp.). If ∂E is inessential in $F''_{D_\alpha D_\beta}$, then ∂E bounds a disk in $F''_{D_\alpha D_\beta}$ containing D'_α and D'_β in its interior. In this case, ∂E is separating in F, which contradicts the assumption of Case 1. If ∂E is essential in $F''_{D_\alpha D_\beta}$, then we get a contradiction similarly as when $\partial E \subset F'_{D_\alpha D_\beta}$ or $F'''_{D_\alpha D_\beta}$.

Case 1-2-3. If $k \geq 1$ and exactly one of $\partial D_0, \cdots, \partial D_m$, say ∂D_α for some α ($0 \leq \alpha \leq m$), is separating in F. Then, $F_{D_\alpha} = F'_{D_\alpha} \cup F''_{D_\alpha}$, $g(F'_{D_\alpha}) = 2$, and $g(F''_{D_\alpha}) = 1$. Here, we get $\partial E \subset F'_{D_\alpha}$ by Lemma 3.1. Therefore, there is some $\partial D_j \subset F'_{D_\alpha}$ for $0 \leq j \neq \alpha \leq m$ by the assumption $k \geq 1$. If we compress F'_{D_α} along E, then $(F'_{D_\alpha})_E$ is a torus. We can assume that there are three disks E', E'', and D'_α in $(F'_{D_\alpha})_E$ where E' (E'', D'_α resp.) is a parallel copy of E (E, D_α resp.) If ∂D_j is inessential in $(F'_{D_\alpha})_E$, then ∂D_j bounds a disk in $(F'_{D_\alpha})_E$ containing (1) E' and E'', (2) E' and D'_α, (3) E'' and D'_α, or (4) E', E'' and D'_α in its interior. In any case, there is a S^2-component in $F_{D_\alpha D_j E}$ determined by D_j and E or D_α, D_j and E. Hence, if we apply Lemma 3.1 to the 1-subsimplex of Δ determined by (D_α, E) and (D_j, E), then we get a contradiction. If ∂D_j is essential in $(F'_{D_\alpha})_E$, then we get a S^2-component in $F_{D_\alpha D_j E}$. Therefore, We get a contradiction similarly.

Case 1-2-4. If $k \geq 1$ and all of $\partial D_0, \cdots, \partial D_m$ are non-separating in F, then we can find two of them ∂D_α and ∂D_β for $0 \leq \alpha \neq \beta \leq m$ such that $\partial D_\alpha \cup \partial D_\beta$ separate F. (Suppose not, i.e. $F_{D_\alpha D_\beta}$ is connected for any choice of α and β for $0 \leq \alpha \neq \beta \leq m$. If ∂E is inessential in $F_{D_\alpha D_\beta}$, then there is a S^2-component in $F_{D_\alpha D_\beta E}$ determined by E and one or both of D_α and D_β. Hence, if we apply Lemma 3.1 to the 1-subsimplex of Δ determined by (D_α, E) and (D_β, E), then we get a contradiction. Hence ∂E is essential in $F_{D_\alpha D_\beta}$. Since $F_{D_\alpha D_\beta}$ is a torus, $F_{D_\alpha D_\beta E}$ is a 2-sphere. Therefore, we get a contradiction similarly.) Now we get $F_{D_\alpha D_\beta} = F'_{D_\alpha D_\beta} \cup F''_{D_\alpha D_\beta}$, where each of $F'_{D_\alpha D_\beta}$ and $F''_{D_\alpha D_\beta}$ is a torus. Without loss of generality, assume that $\partial E \subset F'_{D_\alpha D_\beta}$. If ∂E is inessential in $F'_{D_\alpha D_\beta}$, then there is a S^2-component in $F_{D_\alpha D_\beta E}$ determined by D_α, D_β, and E. Hence, if we apply Lemma 3.1 to the 1-subsimplex of Δ determined by (D_α, E) and (D_β, E), then we get a contradiction. If ∂E is essential in $F'_{D_\alpha D_\beta}$, then this also contradicts to Lemma 3.1 similarly.

Case 2. ∂E is separating in F.

In this case, $F_E = F'_E \cup F''_E$, where $g(F'_E) = 2$ and $g(F''_E) = 1$. By Lemma 3.1, $\partial D_0, \cdots, \partial D_m \subset F'_E$.

Case 2-1. If some ∂D_α for $0 \leq \alpha \leq m$ is separating in F, then $F_{D_\alpha E} = F'_{D_\alpha E} \cup F''_{D_\alpha E} \cup F'''_{D_\alpha E}$ where each component is a torus. Let $F'_{D_\alpha E}$ ($F'''_{D_\alpha E}$ resp.) be the component determined by only D_α (E resp.) and $F''_{D_\alpha E}$ be the component determined by both D_α and E. If $m = 0$, then proof ends. Now we assume that $m \geq 1$. Suppose that there is ∂D_β for $0 \leq \beta \neq \alpha \leq m$ which inessential in $F_{D_\alpha E}$.



FIGURE 3. 1-simplex cases

If $\partial D_\beta \subset F'_{D_\alpha E}$ or $F'''_{D_\alpha E}$, then ∂D_β is isotopic to ∂D_α or ∂E respectively, i.e. a contradiction. Therefore, $\partial D_\beta \subset F''_{D_\alpha E}$. In this case, there is a S^2-component in $F_{D_\alpha D_\beta E}$ determined by D_α, D_β, and E. If we apply Lemma 3.1 to the 1-subsimplex of Δ determined by (D_α, E) and (D_β, E), then we get a contradiction. Therefore, ∂D_β is essential in $F_{D_\alpha E}$ for $0 \leq \beta \neq \alpha \leq m$. Since all components of $F_{D_\alpha E}$ are tori, Lemma 3.1 forces ∂D_β to be in $F'_{D_\alpha E}$. Moreover, we get $m \leq 1$ since ∂D_β is not isotopic to ∂D_α in F and there is only one isotopy class of disjoint essential simple closed curves in $F'_{D_\alpha E}$. In particular, ∂D_β is non-separating in F if $m = 1$.

Case 2-2. If all ∂D_i for $0 \leq i \leq m$ is non-separating in F, then $\partial D_0, \cdots, \partial D_m$ represent different isotopy classes of non-separating curves in F'_E. (Suppose that ∂D_α is isotopic to ∂D_β in F'_E for $0 \leq \alpha \neq \beta \leq m$. Using this isotopy, we get an annulus which ∂D_α and ∂D_β bound in F'_E, i.e. there is a S^2-component in $F_{D_\alpha D_\beta E}$ determined by D_α, D_β, and E. Hence, if we apply Lemma 3.1 to the 1-subsimplex of Δ determined by (D_α, E) and (D_β, E), then we get a contradiction.) If $m = 0$, then the proof ends. If $m \geq 1$, then $(F'_E)_{D_0 D_1}$ must contain a S^2-component determined by D_0, D_1, and E, i.e. $F_{D_0 D_1 E}$ does so. Hence, if we apply Lemma 3.1 to the 1-subsimplex of Δ determined by (D_0, E) and (D_1, E), then we get a contradiction. □

Corollary 3.3. *Assume M, H, and F as in Lemma 3.2. If (D_0, E) and (D_1, E) determine a* 1*-simplex in $\mathcal{WR}$, then ∂D_0 is separating and ∂D_1 is non-separating in F without loss of generality. Moreover, $\partial D_0 \cup \partial D_1$ cuts off a pair of pants from F, and $\tilde{F}$ does not have compressing disks in V other than those parallel to D_0 where $\tilde{F} \subset F$ is obtained from the once-punctured genus two component of $F - \partial D_0$ removing ∂E. We get the same result if (D, E_0) and (D, E_1) determine a* 1*-simplex in $\mathcal{WR}$.*

Proof. In the proof of Lemma 3.2, we have two cases when $\mathcal{WR}$ has a 1-simplex, i.e. $m = 1$, Case 1-2-1 and Case 2-1. In each case, one of ∂D_0 and ∂D_1 is separating and the other is non-separating in F. (In Case 1-2-1, the separating one is ∂D_0 and the non-separating one is ∂D_1. In Case 2-1, the separating one is ∂D_α and the non-separating one is ∂D_β.) Moreover, the separating one cuts off a once-punctured torus from F, and the non-separating one is a non-separating simple closed curve in this torus. Therefore, $\partial D_0 \cup \partial D_1$ cuts off a pair of pants from F (see Figure 3, the left one is when ∂E is non-separating and the right one is when ∂E is separating in F.) Since $\dim(\mathcal{WR}) \leq 1$, the last statement is obvious. □

By Lemma 3.2 we can assume that $\mathcal{WR}$ is a 1-dimensional graph if it is not a set of vertices. Suppose that two vertices $v_1 = (D_1, E_1)$ and $v_2 = (D_2, E_2)$ determine an edge e in $\mathcal{WR}$. Let us label e "D" ("E" resp.) if $E_1 = E_2$ ($D_1 = D_2$ resp.) and call e a *D-edge* (an *E-edge* resp.)

Lemma 3.4. *Assume M, H, and F as in Lemma 3.2. If there are two adjacent edges e and f in $\mathcal{WR}$ such that e is determined by (D_0, E) and (D_1, E) and f is determined by (D_1, E) and (D_2, E), i.e. both edges are labelled "D", then ∂D_1 is non-separating, and ∂D_0, ∂D_2 are separating in F. We get the same result if both edges are labelled E.*

Proof. By Corollary 3.3, we have the following two cases.

Case 1. ∂D_0 and ∂D_2 are non-separating, and ∂D_1 is separating in F.

Case 2. ∂D_0 and ∂D_2 are separating, and ∂D_1 is non-separating in F.

We will show that we only get Case 2. Suppose that Case 1 holds. Let $F_{D_1} = F'_{D_1} \cup F''_{D_1}$, where $g(F'_{D_1}) = 1$ and $g(F''_{D_1}) = 2$. Let V' be the part of V that F'_{D_1} bounds in V. If we consider the edge e, then $\partial D_0 \subset F'_{D_1}$ since $\partial D_0 \cup \partial D_1$ cuts off a pair of pants from F by Corollary 3.3. The existence of the edge e implies $D_0 \cap D_1 = \emptyset$, i.e. $D_0 \subset V'$. Hence, we can compress F'_{D_1} along the disk D_0 in V' and get a 2-sphere S. Since $\partial_- V$ does not have a 2-sphere component, the inside of S in V must be a 3-ball, i.e. V' is a solid torus and D_0 is a meridian disk of V'. If we use Corollary 3.3 for the edge f, then D_2 is also a properly embedded disk in V' and both of D_0 and D_2 must be meridian disks of V'. Therefore, D_0 is isotopic to D_2 in V' and we can assume that this isotopy misses D_1. This means D_0 is isotopic to D_2 in V, which is a contradiction. □

Definition 3.5. If there is a maximal connected subgraph of $\mathcal{WR}$ whose edges are labelled only D and it consists of two or more edges, we call it a *"D-cluster"*. Similarly, we can define an *"E-cluster"*. In a D-cluster (an E-cluster resp.), all vertices have the same E-disk (D-disk resp.).

Lemma 3.6. *A D-cluster does not contain a loop. Moreover, there is only one vertex in a D-cluster which is adjacent to two or more edges in the D-cluster. We get the same result for an E-cluster.*

Proof. If a D-cluster have a subgraph like $v' -_{e_1} v -_{e_2} w -_{e_3} w'$ where e_1 and e_2 share a vertex v and e_2 and e_3 share a vertex w ($v \neq w$), then the boundary of the D-disk of v must be non-separating in F by applying Lemma 3.4 to $v' -_{e_1} v -_{e_2} w$. But if we apply Lemma 3.4 to $v -_{e_2} w -_{e_3} w'$ again, then we get the previous D-disk is separating in F, which is a contradiction. Therefore, a D-cluster cannot have a line segment of length 3 or more. By the definition of $\mathcal{WR}$, there is no loop of length at most two, i.e a D-cluster cannot have a loop.

Suppose that $\bar{v}$ and $\bar{w}$ are different vertices in a D-cluster such that each of both is adjacent to two or more edges in the D-cluster, i.e. there are different two edges e_v^1 and e_v^2 (e_w^1 and e_w^2 resp.) in the D-cluster containing $\bar{v}$ ($\bar{w}$ resp.). Since a D-cluster is connected, there is a path l from $\bar{v}$ to $\bar{w}$ in the D-cluster. We can assume that l does not contain both e_v^1 and e_v^2 since there is no loop in a D-cluster. Similarly, l does not contain both e_w^1 and e_w^2. Without loss of generality, assume that e_v^1 and e_w^1 are not contained in l. If we consider $e_v^1 \cup l \cup e_w^1$, then it is a line segment of length at least three in the D-cluster, i.e. a contradiction. Therefore, there is only one vertex in a D-cluster which is adjacent to two or more edges in the D-cluster. □



FIGURE 4. An example of a D-cluster in $\mathcal{WR}$. (D_0, E) is the center and other vertices are hands.

Definition 3.7. By Lemma 3.6, there is a unique vertex in a D-cluster (an E-cluster resp.) adjacent to two or more edges in the cluster. We call it *"the center of a D-cluster (an E-cluster resp.)"*. We call the other vertices *"hands of a D-cluster (an E-cluster resp.)."* See Figure 4. Note that if an edge $v -_e w$ is contained in a D-cluster (an E-cluster resp.), then one of v and w is the center of the cluster and the other is a hand.

Lemma 3.8. *Assume M, H, and F as in Lemma 3.2 and consider $\mathcal{WR}$. Every edge of $\mathcal{WR}$ is contained in some D-cluster or E-cluster. Moreover, every D- or E- cluster has infinitely many edges.*

Proof. Without loss of generality, assume that there is an edge e labelled D, say $(D_0, E) -_e (D_1, E)$. By Corollary 3.3, we can assume that ∂D_0 is separating and ∂D_1 is non-separating in F. If we compress F along D_1, then F_{D_1} bounds the part of V which misses D_1, say V', in V. Let D_1' and D_1'' be two parallel copies of D_1 on F_{D_1}. Let us consider two disks $\bar{D}_1'$ and $\bar{D}_1''$ such that $\bar{D}_1'$ ($\bar{D}_1''$ resp.) is obtained by extending D_1' (D_1'' resp.) slightly on F_{D_1} and pushing its interior into V' so that it is properly embedded in V' (see Figure 5), and an arc α connecting $\partial\bar{D}_1'$ and $\partial\bar{D}_1''$ on F_{D_1} which misses $\partial\bar{D}_1'$, $\partial\bar{D}_1''$, and ∂E on its interior. If we perform the band sum of $\bar{D}_1'$ and $\bar{D}_1''$ along α in V', then we get a properly embedded disk D_0^α in V' which is ∂-parallel in V'. D_0^α is an essential separating disk in V and we can assume that D_0^α misses D_1 and E. Therefore, we get an edge $(D_0^\alpha, E) -_{e_\alpha} (D_1, E)$ in $\mathcal{WR}$. We claim that there are infinitely many isotopy classes of D_0^α in V. If we compress F_{D_1} again along E, then α is represented by an arc connecting $\partial\bar{D}_1'$ and $\partial\bar{D}_1''$ on F_{D_1E} which misses $\partial\bar{D}_1'$, $\partial\bar{D}_1''$, and the boundary of a parallel copy of E (if ∂E is separating in F_{D_1}) or two parallel copies of E (if ∂E is non-separating in F_{D_1}) on its interior. Note that the genus of each component of F_{D_1E} must be one, otherwise we can find a S^2-component in F_{D_1E} determined by D_1 and E and we get a contradiction by applying Lemma 3.1 to the 0-simplex (D_1, E). Let $\bar{F}_{D_1E}$ be the component of F_{D_1E} containing α. Since $g(\bar{F}_{D_1E}) = 1$, there are infinitely many isotopy classes of such arcs (see Figure 6, we do not fix the endpoints of the arcs on $\partial\bar{D}_1'$ and $\partial\bar{D}_1''$ when we consider these isotopies.) Moreover, if such arcs are non-isotopic, then the resulting disks are also non-isotopic in V. □

Lemma 3.9. *Assume M, H, and F as in Lemma 3.2 and consider $\mathcal{WR}$. If there are consecutive two edges with different labels in $\mathcal{WR}$, then either the boundary*



FIGURE 5. An arc α connecting $\partial\bar{D}_1'$ and $\partial\bar{D}_1''$



FIGURE 6. There are infinitely many isotopy classes of α.

curves of corresponding four disks are all disjoint or the D-cluster and the E-cluster containing the D-edge and the E-edge respectively have the common center. In the latter case, we can replace a separating disk among the four disks by a new separating disk in the same compression body so that the resulting four disks are all disjoint.

Proof. Let $(D_0, E_0) -_e (D_1, E_0) -_f (D_1, E_1)$ be the two edges, where e is labelled D and f is labelled E. By Corollary 3.3, we get a situation like Figure 3 for each edge, but in the case of f, we need to consider one D-disk and two E-disks. Since D_0, D_1, and E_0 are all disjoint, and D_1, E_0, and E_1 are also all disjoint by the existence of the edges e and f, we will prove that $\partial E_1 \cap \partial D_0 = \emptyset$ or the both clusters guaranteed by Lemma 3.8 have the common center. Let the separating one be D_i and the non-separating one be D_j between D_0 and D_1. Let $F_{D_i} = F'_{D_i} \cup F''_{D_i}$, where $g(F'_{D_i}) = 2$ and $g(F''_{D_i}) = 1$. We already know that $\partial E_0 \subset F'_{D_i}$ and $\partial D_j \subset F''_{D_i}$ by Corollary 3.3.

Case 1. $\partial E_1 \cap \partial D_j \neq \emptyset$, i.e. $j = 0$. (so $i = 1$)
In this case, the existence of the vertex (D_1, E_1) implies that $\partial E_1 \cap \partial D_i = \emptyset$. Moreover, $\partial E_1 \subset F''_{D_i}$ from $\partial E_1 \cap \partial D_j \neq \emptyset$. But if ∂E_1 is non-separating in F, then ∂E_1 is essential in F''_{D_i}, i.e. E_1 and D_i determine a S^2 component of $F_{D_i E_1}$. Hence, if we apply Lemma 3.1 to the 0-simplex (D_i, E_1), we get a contradiction. If ∂E_1 is separating in F, then it must be isotopic to ∂D_i, it is also a contradiction.

Case 2. $\partial E_1 \cap \partial D_j = \emptyset$.
If $j = 0$, then the proof ends. Therefore, assume that $j = 1$ (so $i = 0$.) If we apply

Corollary 3.3 to the edge f, then one of ∂E_0 and ∂E_1 is separating and the other is non-separating in F. Let ∂E_k be the separating one and ∂E_l be the non-separating one. Here, ∂D_1 (∂E_l resp.) goes into the genus two component (the genus one component resp.) of F_{E_k} again by Corollary 3.3.

Case 2-1. $\partial E_1 \cap \partial D_i \neq \emptyset$.
Suppose that $l = 1$, (so $k = 0$). Let $F_{E_k} = F'_{E_k} \cup F''_{E_k}$, where $g(F'_{E_k}) = 2$ and $g(F''_{E_k}) = 1$. Here, ∂D_i must go into F'_{E_k} (if $\partial D_i \subset F''_{E_k}$, then it must be parallel to ∂E_k since ∂D_i is separating in F and $g(F''_{E_k}) = 1$.) Since $\partial E_l \cap \partial D_i \neq \emptyset$ from the assumption of Case 2-1, ∂E_l also goes into F'_{E_k}. But we already know that $\partial D_j = \partial D_1$ is contained in F'_{E_k} from the assumption of Case 2. If we compress F'_{E_k} along D_j and E_l, then $(F'_{E_k})_{D_j E_l}$ must have a 2-sphere component determined by D_j, E_k, and E_l since ∂D_j and ∂E_l are two disjoint non-separating simple closed curves in F'_{E_k}. Hence, if we apply Lemma 3.1 to the 1-simpex f, then we get a contradiction. Therefore we get $l = 0$ (so $k = 1$), i.e. both ∂D_1 and ∂E_0 (∂D_0 and ∂E_1 resp.) are non-separating (separating resp.) in F. Let $\mathcal{D}$ ($\mathcal{E}$ resp.) be the D-cluster (the E-cluster resp.) containing the edge e (f resp.) Since the D-disk (E-disk resp.) for the center of a D-cluster (an E-cluster resp.) is non-separating in its compression body by Lemma 3.4, the vertex (D_0, E_0) ((D_1, E_1) resp.) cannot be the center of $\mathcal{D}$ ($\mathcal{E}$ resp.) Therefore, (D_1, E_0) is the common center of both $\mathcal{D}$ and $\mathcal{E}$.

Now we will prove that there is another edge $(D_1, E_0) -_g (D_1, E_2)$ in $\mathcal{E}$ such that $E_2 \cap D_0 = \emptyset$. If we compress F along D_0, then $F_{D_0} = F'_{D_0} \cup F''_{D_0}$ where $g(F'_{D_0}) = 2$ and $g(F''_{D_0}) = 1$. We get $\partial E_0 \subset F'_{D_0}$ and $\partial D_1 \subset F''_{D_0}$ by applying Corollary 3.3 to the edge e. If we compress F'_{D_0} again along E_0, then we get a torus $\bar{F}_{D_0 E_0}$. There are two parallel copies E'_0 and E''_0 of E_0 and one parallel copy D'_0 of D_0 in $\bar{F}_{D_0 E_0}$. If we use the band sum arguments by using an arc connecting E'_0 and E''_0 missing $\partial E'_0$, $\partial E''_0$, and $\partial D'_0$ on its interior as in the proof of 3.8, then we can find an essential separating disk E_2 in W such that $E_2 \cap D_0 = E_2 \cap D_1 = E_2 \cap E_0 = \emptyset$. Therefore, we get an edge $(D_1, E_0) -_g (D_1, E_2)$ which is contained in $\mathcal{E}$. By the existence of the edge g, three disks D_1, E_0, and E_2 are all disjoint. Since we already know that D_0, D_1, and E_0 are all disjoint by the existence of the edge e, $E_2 \cap D_0 = \emptyset$ means that D_0, D_1, E_0, and E_2 are all disjoint.

Case 2-2. $\partial E_1 \cap \partial D_i = \emptyset$.
In this case, $\partial E_1 \cap \partial D_0 = \emptyset$, i.e. the proof ends. □

Corollary 3.10. *Assume M, H, and F as in Lemma 3.2. There are consecutive two edges with different labels, $(\bar{D}, \tilde{E}) -_e (\tilde{D}, \tilde{E}) -_f (\tilde{D}, \bar{E})$ in $\mathcal{WR}$ if and only if the disks $\bar{D}$, $\tilde{D} \subset V$ and $\bar{E}$, $\tilde{E} \subset W$ hold the following conditions.*

1. *Four boundary curves of the disks represent different isotopy classes in F.*
2. *One of $\partial\bar{D}$ and $\partial\tilde{D}$ ($\partial\bar{E}$ and $\partial\tilde{E}$ resp.) is separating and the other is non-separating in F.*
3. *$\partial\bar{D} \cup \partial\tilde{D}$ cuts off a pair of pants from F, and so does $\partial\bar{E} \cup \partial\tilde{E}$. Moreover, both pairs of pants are disjoint in F if the four disks are all disjoint.*
4. *Either (A) the four disks are all disjoint, or (B) we can replace one separating disk among these disks by another separating disk in the same compression body so that the resulting four disks are all disjoint and satisfying the conditions (1), (2), and (3).*



FIGURE 7. An example of the boundaries of four disks satisfying the four conditions of Corollary 3.10

Proof. ($\Leftarrow$) This is obvious since we can assume the four disks $\bar{D}$, $\tilde{D}$, $\bar{E}$, and $\tilde{E}$ are all disjoint.

($\Rightarrow$) Suppose that there are consecutive two edges with different labels, $(\bar{D},\tilde{E})-_e(\tilde{D},\tilde{E})-_f(\tilde{D},\bar{E})$ in $\mathcal{WR}$. If a D-disk and an E-disk share their boundaries, then the splitting is reducible, i.e. the manifold is reducible or the splitting is stabilized, this is a contradiction. Moreover, $\bar{D}$ and $\tilde{D}$ ($\bar{E}$ and $\tilde{E}$ resp.) cannot share their boundaries by the existence of edge e (f resp.). Therefore, we get the condition (1).

If we use Corollary 3.3 for both edges, we get the condition (2).

Suppose that the four disks are all disjoint. Let P_D (P_E resp.) be the pair of pants which $\bar{D}\cup\tilde{D}$ ($\bar{E}\cup\tilde{E}$ resp.) cuts off from F by Corollary 3.3. If $P_D\cap P_E\neq\emptyset$, then one of both belongs to the interior of the other since the boundary curves are disjoint. Assume that $P_E\subset\operatorname{int}(P_D)$ without loss of generality. Then, either some component of ∂P_E is inessential in P_D (so also in F) or each component of ∂P_E is isotopic to some component of ∂P_D in F, any of both cases gives a contradiction. Therefore, we get the condition (3).

If the four disks are all disjoint, then we get the condition (4). Assume that some disk among them intersects the union of the others. This means that Case 2-1 of the proof of Lemma 3.9 holds. Therefore, we can assume that $\bar{D}$, $\tilde{D}$, and $\tilde{E}$ are all disjoint and $\tilde{D}$, $\tilde{E}$, and $\bar{E}$ are all disjoint, but $\bar{E}$ intersects $\bar{D}$. Moreover, we can replace $\bar{E}$ by another separating disk $\bar{E}'$ in W so that we can make an edge $(\tilde{D},\tilde{E})-_g(\tilde{D},\bar{E}')$ and $\bar{D}$, $\tilde{D}$, $\tilde{E}$, and $\bar{E}'$ are all disjoint as in the proof. We can check that this four disks satisfy the conditions (1), (2), and (3) since the edges e and g are also consecutive two edges with different labels. $\square$

We can imagine something like Figure 7 if there exist four disks satisfying the four conditions of Corollary 3.10 and they are all disjoint. The thick curves are separating and the thin curves are non-separating in F.

Now we introduce the key theorem of this article. The idea of this theorem is to isolate one cluster from another cluster in $\mathcal{WR}$.

Theorem 3.11. *Let M be an irreducible 3-manifold and $H=(V,W;F)$ be an unstabilized genus three Heegaard splitting of M. Suppose that there is no choice of four disks $\bar{D}$, $\tilde{D}\subset V$ and $\bar{E}$, $\tilde{E}\subset W$ which satisfy the four conditions of Corollary*

3.10. If we can choose two weak reducing pairs (D_0, E_0) *and* (D_1, E_1) *such that* $\partial D_0 \cap \partial E_1 \neq \emptyset$ *and* $\partial D_1 \cap \partial E_0 \neq \emptyset$ *up to isotopy, then* H *is critical.*

Proof. If (D_0, E_0) and (D_1, E_1) represent the same vertex in $\mathcal{WR}$, then we get $D_0 = D_1$. But $D_0 \cap E_0 = \emptyset$ leads $D_1 \cap E_0 = \emptyset$, which contradicts the assumption of this theorem. Hence, (D_0, E_0) and (D_1, E_1) are different vertices in $\mathcal{WR}$. If $\dim(\mathcal{WR}) = 0$, then the proof ends by Proposition 2.3. Therefore, assume that $\dim(\mathcal{WR}) = 1$ by Lemma 3.2. We will prove that there is no path connecting (D_0, E_0) and (D_1, E_1) in $\mathcal{WR}$ to use Proposition 2.3. If there is a path from (D_0, E_0) to (D_1, E_1) in $\mathcal{WR}$, then each of both vertices belongs to some edge. Hence, Lemma 3.8 forces each of both vertices to be contained in some D- or E-cluster. Now we get the three cases.

Case 1. (D_0, E_0) and (D_1, E_1) belong to the same D-cluster (E-cluster resp.). In this case, $E_0 = E_1$ ($D_0 = D_1$ resp.). But the weak reducing pair (D_0, E_0) means that $\partial D_0 \cap \partial E_0 = \emptyset$, i.e. $\partial D_0 \cap \partial E_1 = \emptyset$ ($\partial D_1 \cap \partial E_0 = \emptyset$ resp.), which contradicts the assumption of this theorem.

Case 2. (D_0, E_0) belongs to some D-cluster (E-cluster resp.) and (D_1, E_1) belongs to another D-cluster (E-cluster resp.)
Suppose that there is a path connecting two vertices, $(D_0, E_0) -_{e_1} - \cdots -_{e_n} (D_1, E_1)$ in $\mathcal{WR}$. This path must contain an edge labelled E (D resp.) otherwise two clusters are united, but this goes to Case 1 and leads a contradiction. Therefore, we can find a line segment of length two in $\mathcal{WR}$ whose edges are labelled differently in the union of the path and these two clusters. This line segment gives the four disks satisfying the four conditions of Corollary 3.10, which contradicts the assumption of this theorem.

Case 3. (D_0, E_0) belongs to some D-cluster (E-cluster resp.) and (D_1, E_1) belongs to some E-cluster (D-cluster resp.).
Suppose that there is a path connecting two vertices. Since two clusters have different labels, we can find a line segment of length two in $\mathcal{WR}$ whose edges are labelled differently in the union of the path and these two clusters, this gives a contradiction as in Case 2. □

Now we will prove Theorem 1.1. Let M, H, F, D_0, D_1, E_0, and E_1 be those in Theorem 1.1.

We divide the proof into the two cases.

Case 1. There is no weak reducing pair whose two disks are separating in their compression bodies.
If there is a choice of four disks satisfying the four conditions of Corollary 3.10, then we can assume that these four disks are all disjoint, i.e. the two separating disks among them correspond to a weak reducing pair, which contradicts the assumption of Case 1. Therefore, there is no choice of four disks satisfying the four conditions of Corollary 3.10. Hence, we can use Theorem 3.11. This completes the proof.

Case 2. There is a weak reducing pair whose two disks are separating in their compression bodies.

We will prove that we cannot choose four disks $\bar{D}$, $\tilde{D} \subset V$, $\bar{E}$, $\tilde{E} \subset W$ satisfying the four conditions of Corollary 3.10.

Suppose that such four disks exist and assume that $\bar{D} \subset V$ and $\bar{E} \subset W$ are separating disks among these four disks. (So $\tilde{D}$ and $\tilde{E}$ are non-separating in their compression bodies.)

Since we can assume $\bar{D}\cap\bar{E}=\emptyset$, $\partial\bar{D}\cup\partial\bar{E}$ cuts F into three pieces F', F'', and F''', where F' (F''' resp.) is the once-punctured torus determined by only $\partial\bar{D}$ ($\partial\bar{E}$ resp.) in F and F'' is the twice-punctured torus determined by both $\partial\bar{D}$ and $\partial\bar{E}$ in F. Since the Heegaard splitting H must be irreducible (otherwise, M is reducible or the splitting is stabilized), we can ignore the possibility that $\partial\bar{D}$ is isotopic to $\partial\bar{E}$ in F. $\bar{D}$ ($\bar{E}$ resp.) cuts off a solid torus or $T^2\times I$ from V (W resp.) where T^2 is a torus. By the assumption of this theorem, we can assume that at least one of $\bar{D}$ and $\bar{E}$, say $\bar{D}$ without loss of generality, cuts off $T^2\times I$ from V, say V'. Let T be the torus component of $\partial_- V$ which realizes the product V'. Our assumption means that V' is the part of V which $F'\cup\bar{D}$ bounds in V.

Since $\partial\bar{D}\cup\partial\tilde{D}$ cuts off a pair of pants from F from Corollary 3.10, $\partial\tilde{D}$ must be a non-separating essential simple closed curve in the interior of F', i.e. $\tilde{D}$ is a compressing disk for $F'=\partial V'\cap\partial_+ V$. Since $\bar{D}$ is a separating disk in V and $\tilde{D}$ is disjoint from $\bar{D}$, we get $\tilde{D}\subset V'$, i.e. F' is compressible in $V'\cong T^2\times I$, this is a contradiction

Therefore, we can use Theorem 3.11. This completes the proof.

4. The proof of Corollary 1.2

In this section, we will prove Corollary 1.2. First, we consider the following lemmas.

Lemma 4.1. *Let M be a closed 3-manifold, H be a genus three Heegaard splitting of M, F be the Heegaard surface. M has a weak reducing pair such that the two disks are separating in their handlebodies and the boundaries of both disks are not isotopic in F if and only if $M=M_1\cup_T M_2$, where each M_i has a genus two Heegaard splitting for $i=1,2$, T is a torus, and H can be represented as an amalgamation of these genus two splittings along T.*

Proof. ($\Leftarrow$) Let the Heegaard splitting of M_1 (M_2 resp.) be $(V_+,W_+;F_1)$ ($(V_-,W_-;F_2)$ resp.), and $W_+\cap V_-=T$. Since a compression body with non-empty minus boundary components can be represented as a union of a product of its minus boundary components and 1-handles, $W_+=(T\times I)\cup(\text{a 1-handle})$ and $V_-=(T\times I)\cup(\text{a 1-handle})$. Let T be the 0-level in $T\times I$ of W_+ and V_- and the attaching disks for the 1-handles go into the 1-levels in these products. In order to proceed the amalgamation, isotope the attaching disks of both 1-handles so that they can be disjoint after the projection into T (see Figure 8.) Note that the projection function may also be changed during this isopoty so that the projection images of these attaching disks do not intersect one another. Let us consider tho curves C_E and C_D in T such that C_E (C_D resp.) bounds a disk in T and this disk contains the projection images of the attaching disks for W_+ (V_- resp.) in its interior. Make sure that the disk for C_E is disjoint from that of C_D. By using the projection function, we can find the corresponding curve $\bar{C}_E$ in $\partial_+ W_+$ ($\bar{C}_D$ in $\partial_+ V_-$ resp.). Consider a disk $E\subset W_+$ ($D\subset V_-$ resp.) such that $\partial E=\bar{C}_E$ ($\partial D=\bar{C}_D$ resp.) like Figure 8. Obviously, D and E are separating in their compression bodies. After amalgamation, D and E are disjoint separating disks in their handlebodies and the boundary of one is not isotopic to that of the other in the resulting Heegaard surface.

($\Rightarrow$) Suppose that there is a weak reducing pair (D,E) where both disks are separating and the boundaries of both disks are not isotopic in F. Let $D\subset V$ and



Figure 8. In the amalgamation case, we can find a weak reducing pair where both disks are separating.

$E \subset W$. Now F_{DE} consists of three tori F'_{DE}, F''_{DE}, and F'''_{DE}, where F'_{DE} (F'''_{DE} resp.) bounds a solid torus V' in V (W' in W resp.). Find a meridian disk D' (E' resp.) in V' (W' resp.) such that D' (E' resp.) is disjoint from D (E resp.). Let T be the surface $F_{D'E'}$. Then, F is obtained from T by attaching boundaries of two 1-handles whose cocores are D' and E' and removing the interiors of attaching disks of these handles. Now we get the right figure of Figure 8. Since $V' \cap W' = \emptyset$, (D', E') is a weak reducing pair. Therefore, if we perform the weak reduction by (D', E'), then we get the generalized Heegaard splitting where both thick surfaces are of genus two and the unique thin surface is torus. □

Lemma 4.2. *Assume M, H, and F as in Lemma 3.2 and consider $\mathcal{WR}$. If there is an edge labelled D (E resp.) in $\mathcal{WR}$ whose E-disk (D-disk resp.) is non-separating in its compression body, then $\mathcal{WR}$ must have a line segment of length two whose edges are labelled differently.*

Proof. Without loss of generality, suppose that there is an edge $(D_0, E) -_e (D_1, E)$ in $\mathcal{WR}$, ∂D_0 is separating, ∂D_1 is non-separating, and ∂E is non-separating in F by Corollary 3.3 and the assumption of this lemma.

Let F'_{D_0} be the genus two component of F_{D_0}. We can check that $\partial E \subset F'_{D_0}$ and $\partial D_1 \cap F'_{D_0} = \emptyset$ by Corollary 3.3. If we compress F'_{D_0} along E again, then $(F'_{D_0})_E$ is a torus. Let E' and E'' be two parallel copies of E_0 and D'_0 be a parallel copy of D_0 in $(F'_{D_0})_E$. If we using the band sum arguments by using an arc connecting E' and E'' missing $\partial E'$, $\partial E''$, and $\partial D'_0$ on its interior as in the proof of Lemma 3.8, we get a family $\{E_\alpha\}$ of infinitely many essential separating disks in W which miss E and D_0. Choose a disk E_α from $\{E_\alpha\}$, and make an edge $(D_0, E_\alpha) -_f (D_0, E)$ labelled E. Therefore, $(D_0, E_\alpha) -_f (D_0, E) -_e (D_1, E)$ is the wanted line segment of length two whose edges are labelled differently. □

Corollary 4.3. *Assume M, H, and F as in Lemma 3.2, consider $\mathcal{WR}$, and add the assumption that M is closed. If $\dim(\mathcal{WR}) = 1$, then $\mathcal{WR}$ must have a line segment of length two whose edges are labelled differently.*

Proof. Without loss of generality, suppose that there is an edge $(D_0, E) -_e (D_1, E)$ in $\mathcal{WR}$, ∂D_0 is separating, and ∂D_1 is non-separating in F by Corollary 3.3. If

∂E is non-separating in F, then the proof ends by Lemma 4.2. Therefore, assume that ∂E is separating in F, i.e. E is separating in W. Hence, E cuts W into a solid torus W' and a genus two handlebody, where W' is bounded by $\tilde{F} \cup E$ in W. ($\tilde{F}$ is a once-punctured torus which ∂E cut off from F.) Let E_m be a meridian disk of W'. Isotope E_m in W' so that ∂E_m is contained in the interior of $\tilde{F}$. Let $\bar{F}$ be the once-punctured genus two component of $F - \partial D_0$. Since $\partial E \subset \bar{F}$ by Corollary 3.3 and ∂E cuts $\bar{F}$ into a twice-punctured torus and a once-punctured torus (otherwise, ∂E is isotopic to ∂D_0 in $\bar{F}$), we get either $\tilde{F} \subset \bar{F}$ or $\tilde{F}$ is of genus two. The latter case contradicts that $\tilde{F}$ is a once-punctured torus. Hence, $\partial D_0 \cap \partial E_m = \emptyset$, i.e. we get an edge $(D_0, E_m) -_f (D_0, E)$ labelled E. Therefore, $(D_0, E_m) -_f (D_0, E) -_e (D_1, E)$ is the wanted line segment of length two whose edges are labelled differently. □

Let us prove Corollary 1.2. If M is not an amalgamation of two genus two splitting along a torus, then we cannot choose a weak reducing pair whose disks are separating in their hadlebodies by Lemma 4.1. (Since M is irreducible and the splitting is unstabilized, the splitting is irreducible, i.e. we can get rid of the possibilities of weak reducing pairs whose disks have isotopic boundaries.) Hence, there is no weak reducing pair such that each of both disks cuts off a solid torus in its compression body since the genus of the splitting is three. Therefore, if we use Theorem 1.1, then the proof ends.

In addition, we can induce some equivalent conditions for $\dim(\mathcal{WR}) = 0$ when M is closed.

Corollary 4.4. *Let M be a closed orientable irreducible 3-manifold and $H = (V, W; F)$ be an unstabilized weakly reducible genus three Heegaard splitting. Then, the following three statements are equivalent.*

1. *There is no weak reducing pair such that each of both disks cuts off a solid torus in its handlebody.*
2. *H cannot be represented as an amalgamation of genus two splittings along a torus.*
3. $\dim(\mathcal{WR}) = 0$.

Proof. Since M is irreducible, (1) and (2) are equivalent by Lemma 4.1. Therefore we will prove that (1) and (3) are equivalent.

(1)⇒(3) Suppose that $\dim(\mathcal{WR}) = 1$, i.e. the negation of $\dim(\mathcal{WR}) = 0$ by Lemma 3.2. Then, $\mathcal{WR}$ must have a line segment of length two whose edges are labelled differently by Corollary 4.3. This gives the four disks $\bar{D}$, $\tilde{D} \subset V$ and $\bar{E}$, $\tilde{E} \subset W$ satisfying the four conditions of Corollary 3.10. Let $\bar{D}$ and $\bar{E}$ be the two separating disks among them. Since we can assume $\bar{D} \cap \bar{E} = \emptyset$, we get a weak reducing pair of separating disks $(\bar{D}, \bar{E})$. By the assumption that M is closed, $\bar{D}$ ($\bar{E}$ resp.) cuts off a solid torus from V (W resp.).

(3)⇒(1) Suppose that $\dim(\mathcal{WR}) = 0$. Suppose that there is weak reducing pair (D, E) such that each of both disks cuts off a solid torus in its handlebody. Let the solid torus which D cuts off from V be V' and D_m be a meridian disk of V'. If we isotope D_m in V' so that ∂D_m misses D, then we get a 1-simplex $(D, E) - (D_m, E)$ in $\mathcal{WR}$, i.e $\dim(\mathcal{WR}) = 1$ and this gives a contradiction. Therefore, There is no weak reducing pair such that each of both disks cuts off a solid torus in its handlebody. □



FIGURE 9. The standard Heegaard splitting of genus three of T^3, and its compressing disks which realize the critical Heegaard splitting.

5. EXAMPLES OF THEOREM 1.1

5.1. **The three-torus T^3.** It is well known that $T^3 =$ (torus) $\times S^1$ has the unique minimal genus three Heegaard splitting (see Theorem 4.2 of [5] and Theorem 5.7 of [11].)

Let C be a cube $\{(x, y, z) \in \mathbb{R}^3 | -1 \leq x, y, z \leq 1\}$ (see (a) of Figure 9.) If we identify the three pairs of opposite faces of C, we get a 3-torus $M =$ (torus) $\times S^1$. Let $q : C \to M$ be the quotient map. An image by q of a tubular neighborhood of union of three axis in C is a genus three handlebody V. Let $\overline{M - V}$ be W. Then it is easy to see that W is also a genus three handlebody. This is the standard Heegaard splitting of genus three for T^3.

We will prove that this splitting is critical by using Theorem 1.1. If there is a weak reducing pair for the standard splitting of genus three for T^3, then both disks are non-separating in their handlebodies (see Lemma 6 of [6].) Therefore, there is no choice of a weak reducing pair such that each of both disks cuts off a solid torus in its handlebody. By the existence of the weak reducing pairs (D_0, E_0) and (D_1, E_1) as in Figure 9, we can induce that this splitting is critical by Theorem 1.1. This is an example of $\dim(\mathcal{WR}) = 0$ (see Corollary 4.4.) Also we can check that the standard splitting of genus three for T^3 cannot be represented by an amalgamation of two genus two Heegaard splittings along a torus by Corollary 4.4.

Let us check Corollary 1.3 for this example. Let S be the horizontal torus in T^3 in the left of Figure 10 and F be the Heegaard surface of this splitting. Since the fibers of the torus bundle $T^3 = T^2 \times S^1$ is incompressible in T^3, S is incompressible. Here, we can check that $S \cap F$ is inessential in S. Let c be an essential closed curve in S as in the left of Figure 10. If we push S upward along c until we get the right of Figure 10, then we get a torus S' isotopic to S. $S' \cap F$ consists of two circles, where each of both is depicted as a dotted curve in the right of Figure 10. These curves are all essential in both F and S'. Therefore, S' is the surface satisfying Corollary 1.3.

Note that there is another proof to show that this splitting is critical (see [7].)



FIGURE 10. S' is satisfying Corollary 1.3.



FIGURE 11. A tunnel system of the three component chain

5.2. **The three component chain exterior.** Let M be the exterior of the three component chain pictured in Figure 11. (The manifold M may also be considered as $P \times S^1$, where P is a pair of pants.) Let T_1, T_2, and T_3 be the three boundary components of ∂M, and let us consider two arcs a_1 and a_2 such that T_1 and T_2 (T_3 and T_2 resp.) are connected by a_1 (a_2 resp.) as in Figure 11. It is easy to show that $\{a_1, a_2\}$ is a tunnel system for M. Therefore we get a Heegaard splitting $(V, W; F)$ of M, where V is a genus three compression body whose minus boundary consists of T_1, T_2, and T_3 and W is a genus three handlebody. If the splitting is stabilized, then we can reduce the tunnel system. But a compression body whose minus boundary consists of three tori must be of at least genus three, we cannot reduce this tunnel system, i.e. the splitting is unstabilized. Let D_0 (D_1 resp.) be the cocore of a_1 (a_2 resp.) and E_0 (E_1 resp.) be the compressing disk for W depicted as in Figure 11. Here, (D_0, E_0) and (D_1, E_1) are weak reducing pairs. Moreover, $D_0 \cap E_1 \neq \emptyset$ and $D_1 \cap E_0 \neq \emptyset$. If there is an essential separating disk in V, then it cannot cut off a compression body with empty minus boundary from V. That is, there is no weak reducing pair such that each of both disks cuts off a solid torus in its compression body. Therefore, this splitting is critical by Theorem 1.1.

Now we claim that $\dim(\mathcal{WR}) = 1$. Let $\bar{E}_0$ and $\tilde{E}_0$ be two parallel copies of E_0 as in Figure 12. If we consider the band sum of $\bar{E}_0$ and $\tilde{E}_0$ by the arc α connecting $\bar{E}_0$ and $\tilde{E}_0$ as in Figure 12, then we get an essential separating disk E' in W. Since $E' \cap D_0 = \emptyset$, $(D_0, E_0) - (D_0, E')$ is a 1-simplex in $\mathcal{WR}$. This example means that



FIGURE 12. We can find the disk E' by the band sum of two parallel copies of E_0

the first statement and the third statement of Corollary 4.4 are not equivalent for manifolds with non-empty boundary.

Note that E. Sedgwick proved that this splitting is not of minimal genus in [13].

ACKNOWLEDGMENTS

The author is grateful to Prof. Jung-Hoon Lee for carefully reading the paper. The author was supported by the National Research Foundation of Korea grant funded by the Korean Government(Ministry of Education, Science and Technology)[NRF-2011-355-C00014].

REFERENCES

[1] D. Bachman, *Critical Heegaard surfaces.* Trans. Amer. Math. Soc. **354** (2002), no. 10, 4015–4042 (electronic).

[2] D. Bachman, *Connected sums of unstabilized Heegaard splittings are unstabilized*, Geom. Topol. **12** (2008), no. 4, 2327–2378.

[3] D. Bachman, *Topological index theory for surfaces in 3-manifolds*, Geom. Topol. **14** (2010), no. 1, 585–609.

[4] A. J. Casson and C. McA. Gordon, *Reducing Heegaard splittings*, Topology Appl. **27** (1987), no. 3, 275–283.

[5] C. Frohman and J. Hass, *Unstable minimal surfaces and Heegaard splittings*, Invent. Math. **95** (1989), no. 3, 529–540.

[6] J. Johnson, *Automorphisms of the three-torus preserving a genus-three heegaard splitting*, Pacific J. Math. **253** (2011), no. 1, 75–94.

[7] J. Lee, *Examples of unstabilized critical Heegaard surfaces*, arXiv:1005.0974.

[8] Y. Moriah, *On boundary primitive manifolds and a theorem of Casson-Gordon*, Topology Appl. **125** (2002), no. 3, 571–579.

[9] M. Saito, M. Scharlemann and J. Schultens, *Lecture notes on generalized Heegaard splittings*, arXiv:math/0504167v1

[10] M. Scharlemann and J. Schultens, *The tunnel number of the sum of n knots is at least n*, Topology **38** (1999), no. 2, 265–270.

[11] J. Schultens, *The classification of Heegaard splittings for (compact orientable surface)$\times S^1$*, Proc. London Math. Soc. (3) **67** (1993), no. 2, 425–448.

[12] J. Schultens, *Additivity of tunnel number for small knots*, Comment. Math. Helv. **75** (2000), no. 3, 353–367.

[13] E. Sedgwick, *Genus two 3-manifolds are built from handle number one pieces*, Algebr. Geom. Topol. **1** (2001), 763–790.

[14] A. Thompson, *The disjoint curve property and genus 2 manifolds*, Topology Appl. **97** (1999) 273–279.

[15] F. Waldhausen, *Heegaard-Zerlegungen der 3-sphäre*, Topology, **7** (1968), 195–203.

Department of Mathematics, Chung-Ang University, Seoul, Korea
E-mail address: `pibonazi@gmail.com`